

Mindful Continuing Education

Depression in New Mothers

1. The baby blues are experienced by what percentage of women?

- A. 50-60 %
 - B. 60-70%
 - C. 70-80%
 - D. 80-90%
-

2. According to the author, risk factors for postpartum depression may include each of the following EXCEPT:

- A. Experiencing postpartum depression in a previous pregnancy, a family history of depression or mood disorders or a stressful life
 - B. Pregnancy complications, a baby born with special needs or health, or twins, triplets, or multiples born
 - C. A diagnosis of an anxiety disorder, a personality disorder, or a co-occurring mental health/substance abuse issue
 - D. Having issues in your relationship, having a compromised immune system, or experiencing financial hardship
-

3. The baby blues is equally as dangerous as postpartum depression for most women.

- A. True
 - B. False
-

4. Which attachment style do people strive for with their children?

- A. Secure
 - B. Avoidant
 - C. Anxious
 - D. Disorganized
-

5. A multidisciplinary treatment team involves which of the following providers?

- A. Therapist
 - B. Massage therapist
 - C. Psychiatrist
 - D. Physician
 - E. Any professional above and any who will benefit the ongoing health of the mother
-

6. Women with depression are more likely to misuse substances?

- A. True**
 - B. False**
-

7. Interpersonal therapy, which has been found to be clinically beneficial for women with postpartum depression, explores four different interpersonal problems, including role transition, role dispute, interpersonal deficits, and:

- A. Grief**
 - B. Unwanted thoughts and feelings**
 - C. Lack of personal acceptance**
 - D. Relationship issues**
-

8. Of the following treatment services, what is most likely one that should be started right away for women with postpartum depression?

- A. Medication**
 - B. Group therapy**
 - C. Family therapy**
 - D. Individual therapy**
-

9. What should be completed as soon as possible once a provider begins working with a new mother with depression?

- A. A long term plan for what recovery looks like**
 - B. Safety plan**
 - C. Goal setting**
 - D. Cope ahead plan**
-

10. The diagnosis of postpartum depression will involve a screening tool for depression.

- A. True**
 - B. False**
-